

Understanding Peripheral Edema and Long-Term Management

Peripheral edema is swelling caused by the accumulation of excess fluid in the tissues, most commonly affecting the feet, ankles, legs, and sometimes the hands. It is a symptom rather than a disease and can result from a variety of medical conditions.



Common Causes



Heart Disease
Heart failure can cause fluid buildup in the legs, ankles, abdomen, and lungs.



Kidney Disease
Impaired kidney function may lead to sodium and water retention.



Liver Disease
Conditions such as cirrhosis reduce protein production and increase fluid accumulation.



Venous Insufficiency
Weak or damaged leg veins allow fluid to leak into tissues.



Lymphedema
Blockage or dysfunction of lymphatic drainage. Swelling is often persistent.



Medications
Certain medications (e.g., amlodipine, NSAIDs, steroids, hormonal therapies).



Pregnancy
Mild swelling is common, especially in the third trimester.



Prolonged Sitting or Standing
Common during travel, office work, or limited mobility.

Symptoms



Swelling of feet, ankles, or legs



Tight or shiny skin



Indentation after pressing on the skin (pitting edema)



Heaviness or discomfort in the legs



Reduced flexibility of joints



Weight gain from fluid retention



Warning Signs – Seek Immediate Medical Attention

- Shortness of breath
- Chest pain
- Rapid weight gain
- One-sided leg swelling with pain or redness
- Fainting or severe dizziness

These symptoms may indicate serious conditions such as:

- Deep vein thrombosis (DVT)
- Pulmonary embolism
- Acute heart failure

Evaluation



Physical Examination
Assessment of swelling and underlying causes



Laboratory Testing
Kidney & liver function, electrolytes, urinalysis, protein levels



Imaging Studies
Venous ultrasound, echocardiogram, chest X-ray, and other studies when indicated

Treatment and Management

Treatment focuses on the underlying cause and includes lifestyle measures, medications, and long-term follow-up.

Lifestyle Measures



Elevate legs above heart level several times daily



Regular exercise (e.g., walking)



Compression stockings when appropriate



Maintain a healthy weight



Reduce sodium intake



Avoid prolonged sitting or standing

Medications



- Diuretics ("water pills") for fluid overload
- Medications to improve heart function
- Treatment of kidney or liver disease
- Anticoagulants if a blood clot is present



Medication-related edema often improves after adjusting or changing the medication.

Long-Term Management



Regular follow-up based on the underlying condition

- Control blood pressure
- Monitor weight and symptoms
- Use compression therapy when appropriate
- Maintain a healthy weight and active lifestyle
- Manage kidney, liver, heart, or lymphatic disease

Activity Recommendations

Generally Encouraged



Walking



Swimming



Cycling



Light resistance training



Avoid



Prolonged immobility



Excessive dietary sodium



Tight clothing that restricts circulation

Preventing Complications



Inspect skin daily



Prevent skin injury



Treat cuts promptly



Wear properly fitting shoes



Maintain good hydration



Stay physically active

Potential Complications

- Skin breakdown
- Cellulitis
- Venous ulcers
- Reduced mobility
- Recurrent infections

Prognosis

- Depends on the underlying cause.
- Venous insufficiency often improves with compression and exercise.
- Heart, kidney, or liver disease may require ongoing management.
- Lymphedema is usually chronic but can be effectively controlled.



Dr. Shabib Alhadheri, MD
Advanced Pediatric Cardiology

1478 W Elliot Rd, Suite 102, Gilbert, AZ 85233 480-855-1339 www.apcario.com